



## Macroalgal protein hydrolysates from *Palmaria palmata* influence the 'incretin effect' in vitro via DPP-4 inhibition and upregulation of insulin, GLP-1 and GIP secretion

McLaughlin, C., Harnedy-Rothwell, P. A., Lafferty, R., Sharkey, S., V, P., Allsopp, P. J., McSorley, E. M., Fitzgerald, R., & O'Harte, F. (2021). Macroalgal protein hydrolysates from *Palmaria palmata* influence the 'incretin effect' in vitro via DPP-4 inhibition and upregulation of insulin, GLP-1 and GIP secretion. *European Journal of Nutrition*, 60(8), 4439-4452. <https://doi.org/10.1007/s00394-021-02583-3>

[Link to publication record in Ulster University Research Portal](#)

**Published in:**  
European Journal of Nutrition

**Publication Status:**  
Published (in print/issue): 01/12/2021

**DOI:**  
[10.1007/s00394-021-02583-3](https://doi.org/10.1007/s00394-021-02583-3)

**Document Version**  
Author Accepted version

**General rights**  
Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**  
The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk).

# European Journal of Nutrition

## Macroalgal protein hydrolysates from *Palmaria palmata* influence the ‘incretin effect’ in vitro via DPP-4 inhibition and upregulation of insulin, GLP-1 and GIP secretion --Manuscript Draft--

<b>Manuscript Number:</b>	EJON-D-20-01386R2	
<b>Full Title:</b>	Macroalgal protein hydrolysates from <i>Palmaria palmata</i> influence the ‘incretin effect’ in vitro via DPP-4 inhibition and upregulation of insulin, GLP-1 and GIP secretion	
<b>Article Type:</b>	Original Contribution	
<b>Keywords:</b>	Antidiabetic; Dipeptidylpeptidase-4; Dulse; Incretin secretion; <i>Palmaria palmata</i> ; Protein hydrolysate; Type 2 diabetes	
<b>Corresponding Author:</b>	Finbarr O'Harte, PhD Ulster University - Coleraine Campus: Ulster University Coleraine, Co. Londonderry UNITED KINGDOM	
<b>Corresponding Author Secondary Information:</b>		
<b>Corresponding Author's Institution:</b>	Ulster University - Coleraine Campus: Ulster University	
<b>Corresponding Author's Secondary Institution:</b>		
<b>First Author:</b>	Christopher M McLaughlin, BSc PhD	
<b>First Author Secondary Information:</b>		
<b>Order of Authors:</b>	Christopher M McLaughlin, BSc PhD	
	Padraigin A Harnedy-Rothwell, BSc PhD	
	Ryan A Lafferty, BSc PhD	
	Shaun Sharkey, BSc MSc	
	Vadivel Parthasarathy, BSc MSc PhD	
	Philip J Allsopp, BSc PhD	
	Emeir M McSorley, BSc PhD	
	Richard J FitzGerald, BSc PhD	
	Finbarr P M O'Harte, PhD	
<b>Order of Authors Secondary Information:</b>		
<b>Funding Information:</b>	Department of Agriculture, Food and the Marine (IE) (11/F/063 and 11/F064 and 13/F536 and 14/F873)	Professor Richard J FitzGerald
	Department of Education & Learning, Northern Ireland (PhD Studentship)	Dr Christopher M McLaughlin
<b>Abstract:</b>	<p><b>Purpose</b> This study investigated metabolic benefits of protein hydrolysates from the macroalgae <i>Palmaria palmata</i>, previously shown to inhibit dipeptidylpeptidase-4 (DPP-4) activity in vitro.</p> <p><b>Methods</b> Previously, Alcalase/Flavourzyme-produced <i>P. palmata</i> protein hydrolysate (PPPH) improved glycaemia and insulin production in streptozotocin-induced diabetic mice. Here the PPPH, was compared to alternative Alcalase, bromelain and Promod derived hydrolysates and an unhydrolysed control. All PPPH's underwent simulated gastrointestinal digestion (SGID) to establish oral bioavailability. PPPH's and their SGID counterparts were tested in pancreatic, clonal BRIN-BD11 cells to assess their insulinotropic effect and associated intracellular mechanisms. PPPH actions on the</p>	

	<p>incretin effect were assessed via measurement of DPP-4 activity, coupled with GLP-1 and GIP release from GLUTag and STC-1 cells, respectively. Acute in vivo effects of Alcalase/Flavourzyme PPPH administration on glucose tolerance and satiety were assessed in overnight-fasted mice.</p> <p><b>Results</b></p> <p>PPPH's (0.02-2.5 mg/ml) elicited varying insulinotropic effects (<math>p &lt; 0.05</math>–<math>0.001</math>). SGID of the unhydrolysed protein control, bromelain and Promod PPPH's retained, or improved, bioactivity regarding insulin secretion, DPP-4 inhibition and GIP release. Insulinotropic effects were retained for all SGID-hydrolysates at higher PPPH concentrations. DPP-4 inhibitory effects were confirmed for all PPPH's and SGID counterparts (<math>p &lt; 0.05</math>–<math>0.001</math>). PPPH's were shown to directly influence the incretin effect via upregulated GLP-1 and GIP (<math>p &lt; 0.01</math>–<math>0.001</math>) secretion in vitro, largely retained after SGID. Alcalase/Flavourzyme PPPH produced the greatest elevation in cAMP (<math>p &lt; 0.001</math>, 1.7-fold), which was fully retained post-SGID. This hydrolysate elicited elevations in intracellular calcium (<math>p &lt; 0.01</math>) and membrane potential (<math>p &lt; 0.001</math>). In acute in vivo settings, Alcalase/Flavourzyme PPPH improved glucose tolerance (<math>p &lt; 0.01</math>–<math>0.001</math>) and satiety (<math>p &lt; 0.05</math>–<math>0.001</math>).</p> <p><b>Conclusion</b></p> <p>Bioavailable PPPH peptides may be useful for the management of T2DM and obesity.</p>
<b>Suggested Reviewers:</b>	<p>Dagmar Stengel, PhD Botany and Plant Science, NUIG: National University of Ireland Galway dagmar.stengel@nuigalway.ie Has expertise in seaweed and plant sciences and extraction of components from <i>Palmaria palmata</i> seaweed.</p> <p>Chao Zhao, PhD Agriculture Forestry and Fisheries Research Council Secretariat Tsukuba Office Department of Research and Information: Norin Suisan Kenkyu Joho Sogo Center zhchao@live.cn Has expertise in antidiabetic effects of seaweed in diabetic animal models</p> <p>Yoav D Livney, PhD Professor, Technion Israel Institute of Technology livney@technion.ac.il Expertise in proteins from macroalgae for food applications</p> <p>Elis Lima, PhD University of the Azores Department of Agricultural Sciences: Universidade dos Acores Departamento de Ciencias Agrarias elis@uac.pt This author has recently published articles on the bioactivity of macroalgae.</p> <p>Yang Ruijin, PhD Jiangnan University School of Food Science and Technology yrj@jiangnan.edu.cn These authors have extensive experience in isolation of proteins and hydrolysates from seaweeds and have recently published in this same area.</p>

<sup>1</sup>McLaughlin, C. M.; <sup>2,3</sup>Harnedy-Rothwell, P. A.; <sup>1</sup>Lafferty, R. A.; <sup>1</sup>Sharkey, S.; <sup>1</sup>Parthsarathy, V.; <sup>3</sup>Allsopp, P. J.; <sup>3</sup>McSorley, E. M.; <sup>2,3</sup>FitzGerald, R. J. and <sup>1</sup>O'Harte, F.P.M.

<sup>1</sup>School of Biomedical Sciences, Ulster University, Cromore Road, Coleraine, Co. Derry, BT52 1SA, Northern Ireland.

<sup>2</sup>Department of Biological Sciences, University of Limerick, Castletroy, Limerick, Ireland.

<sup>3</sup>Health Research Institute (HRI), University of Limerick, Limerick, Ireland.<sup>4</sup>Nutrition Innovation Centre for Food and Health, School of Biomedical Sciences, Ulster University, Cromore Road, Coleraine, Co. Derry, BT52 1SA, Northern Ireland.

**Key words:** Antidiabetic, Dipeptidylpeptidase-4, Dulce, Incretin secretion, *Palmaria palmata*, Protein hydrolysate, Type 2 diabetes.

**Character count:** 49,519 (Including abstract, references, tables and legends)

### **Corresponding author**

Prof. Finbarr O'Harte  
School of Biomedical Sciences  
Ulster University  
Coleraine, BT52 1SA  
Co. Derry  
Northern Ireland.

**T:** [+44 (0)28 70 124853]

**E:** [fpm.oharte@ulster.ac.uk]

**W:** [www.ulster.ac.uk/](http://www.ulster.ac.uk/)

**W:** <https://www.ulster.ac.uk/faculties/life-and-health-sciences/schools/biomedical-sciences>

## Abstract

### Purpose

This study investigated metabolic benefits of protein hydrolysates from the macroalgae *Palmaria palmata*, previously shown to inhibit dipeptidylpeptidase-4 (DPP-4) activity *in vitro*.

### Methods

Previously, Alcalase/Flavourzyme-produced *P. palmata* protein hydrolysate (PPPH) improved glycaemia and insulin production in streptozotocin-induced diabetic mice. Here the PPPH, was compared to alternative Alcalase, bromelain and Promod derived hydrolysates and an unhydrolysed control. All PPPH's underwent simulated gastrointestinal digestion (SGID) to establish oral bioavailability. PPPH's and their SGID counterparts were tested in pancreatic, clonal BRIN-BD11 cells to assess their insulintropic effect and associated intracellular mechanisms. PPPH actions on the incretin effect were assessed via measurement of DPP-4 activity, coupled with GLP-1 and GIP release from GLUTag and STC-1 cells, respectively. Acute *in vivo* effects of Alcalase/Flavourzyme PPPH administration on glucose tolerance and satiety were assessed in overnight-fasted mice.

### Results

PPPH's (0.02-2.5 mg/ml) elicited varying insulintropic effects ( $p < 0.05$ –0.001). SGID of the unhydrolysed protein control, bromelain and Promod PPPH's retained, or improved, bioactivity regarding insulin secretion, DPP-4 inhibition and GIP release. Insulintropic effects were retained for all SGID-hydrolysates at higher PPPH concentrations. DPP-4 inhibitory effects were confirmed for all PPPH's and SGID counterparts ( $p < 0.05$ –0.001). PPPH's were shown to directly influence the incretin effect via upregulated GLP-1 and GIP ( $p < 0.01$ –0.001) secretion *in vitro*, largely retained after SGID. Alcalase/Flavourzyme PPPH produced the

greatest elevation in cAMP ( $p < 0.001$ , 1.7-fold), which was fully retained post-SGID. This hydrolysate elicited elevations in intracellular calcium ( $p < 0.01$ ) and membrane potential ( $p < 0.001$ ). In acute *in vivo* settings, Alcalase/Flavourzyme PPPH improved glucose tolerance ( $p < 0.01$ – $0.001$ ) and satiety ( $p < 0.05$ – $0.001$ ).

## Conclusion

Bioavailable PPPH peptides may be useful for the management of T2DM and obesity.

## 1    **Introduction**

2    Type 2 diabetes mellitus (T2DM) is a metabolic disorder of complex aetiology characterised  
3    by a deficiency, and/or dysfunction of endogenous insulin and glucagon production [1]. In  
4    diabetes, loss of insulin-producing pancreatic beta-cell mass [2], is accompanied by  
5    dysfunction of glucagon-producing alpha-cells, which fail to respond to the normal suppressive  
6    effects of glucose and insulin, leading to hyperglucagonaemia [3]. Best-estimates state that  
7    there are around 463 million adults globally living with diabetes, projected to rise to 700  
8    million by 2045 [4], with T2DM representing ~90% of cases. The economic burden of diabetes  
9    on global healthcare systems is considerable, with a minimum of \$760 billion USD attributed  
10    to spending on the disease in 2019, equating to 10% of global adult healthcare costs [4]. An  
11    important factor in this spending arises from costs amassed from treatment of microvascular  
12    (retinopathy, nephropathy, and neuropathy) and macrovascular (coronary artery disease,  
13    stroke, and peripheral vascular disease) complications [5]. Preventative strategies, coupled  
14    with earlier diagnosis and novel treatments, have the potential to reduce the occurrence of these  
15    complications [6].

16        Lifestyle interventions such as increased physical activity and improved, nutritionally  
17    balanced diets are considered first-line options in the prevention and treatment of T2DM [7,8].  
18    While high-quality dietary protein is an integral part of any such diet [9], it has also been  
19    established that a high-protein diet can lower postprandial blood glucose in T2DM and improve  
20    overall glucose and lipid metabolism [10,11]. Beyond dietary protein, protein hydrolysates,  
21    peptides and single amino acids can beneficially regulate glycaemia, with the magnitude of  
22    response differing significantly depending on the primary sequence of peptides and specific  
23    amino acids generated following digestion [12,13].

24        Mechanisms determining glycaemic improvements of various protein hydrolysates have  
25    been established, highlighting the importance of inhibitory actions on the ubiquitous enzyme

1 dipeptidylpeptidase-4 (DPP-4) [14-18]. DPP-4 inhibition has become a staple of diabetes  
2 management, with a plethora of drugs now available since the approval of sitagliptin (Januvia®)  
3 in 2006 [19]. Success of DPP-4 inhibition lies in the preservation of the “incretin effect”, which  
4 promotes a rise in plasma insulin following food intake [20]. The rise in plasma insulin not  
5 only reflects a response to increased postprandial glucose, but approximately 50% of the  
6 overall insulinotropic response is attributed to the release of two gut-derived hormones,  
7 namely: glucagon-like peptide 1 (GLP-1) and glucose dependent insulinotropic polypeptide  
8 (GIP) [20]. Both GLP-1 and GIP are inactivated following N-terminal dipeptide removal by  
9 DPP-4 [19]. Furthermore, DPP-4 resistant GLP-1 receptor agonists (incretin mimetics) have  
10 also been developed and are widely prescribed for T2DM management [21-23].

11 It has recently been uncovered that, beyond DPP-4 inhibition, protein hydrolysates from  
12 underutilised marine sources, such as blue whiting, boarfish and salmon skin, can directly  
13 influence glycaemia through improved insulin production and secretion coupled with  
14 upregulated GLP-1 secretion in both *in vitro* [24-26] and *in vivo* settings [25,27]. The present  
15 study has sought to establish whether crude hydrolysates of the macroalgae *Palmaria palmata*  
16 can replicate the effects of piscine-derived protein hydrolysates.

17 *P. palmata* (dulse) has become popular as a foodstuff due to its relatively high protein  
18 content [28], in addition to being a potential source of biofunctional proteinaceous and  
19 antioxidant ingredients [29-31]. Notably, both crude hydrolysates of *P. palmata* [32] and  
20 isolated peptides from this source have demonstrated an ability to inhibit DPP-4 *in vitro* [33].  
21 Furthermore, twice daily, chronic administration of a crude *P. palmata* protein hydrolysate,  
22 Alcalase/Flavourzyme PPPH, has been shown to improve glycaemic control in streptozotocin-  
23 induced diabetic mice [34]. Thus, the present study aims to employ a number of established  
24 screening methods to uncover the specific mechanisms responsible for the positive glycaemic  
25 effects of PPPH's and identify the hydrolysate which shows greatest anti-diabetic potential.



## 1    **Materials and Methods**

### 2    **Materials and chemicals**

3    H-Gly-Pro-AMC (7-amino-4-methyl coumarin) and Diprotin A were obtained from Bachem  
4    Feinchemikalien (Bubendorf, Switzerland). Promod 144 MG provided by Biocatalysts Ltd.  
5    (Cardiff, Wales, UK). HPLC grade water and acetonitrile from VWR International (Dublin,  
6    Ireland) and trinitrobenzenesulphonic acid (TNBS) reagent was from Medical Supply Co Ltd.  
7    (Dublin, Ireland). Calcium chloride dihydrate ( $\text{CaCl}_2 \times 2\text{H}_2\text{O}$ ), D-glucose, HEPES,  
8    hydrochloric acid (HCl), magnesium sulphate ( $\text{MgSO}_4 \times 7\text{H}_2\text{O}$ ), potassium dihydrogen  
9    orthophosphate ( $\text{KH}_2\text{PO}_4$ ), potassium chloride (KCl), sodium bicarbonate ( $\text{NaHCO}_3$ ) and  
10    sodium chloride (NaCl) were purchased from BDH Chemicals Ltd. (Poole, Dorset, UK). Foetal  
11    bovine serum (FBS), Hank's buffered saline solution (HBSS 10x stock), penicillin–  
12    streptomycin (0.1 g/l), RPMI-1640 culture media, Dulbecco's modified Eagle's medium  
13    (DMEM) containing high glucose and trypsin/EDTA (10x) were obtained from Gibco Life  
14    Technologies Ltd. (Paisley, Strathclyde, UK). Radio-labelled sodium iodide ( $\text{Na}^{125}\text{I}$ , IMS 100  
15    mCi/ml stock) was from Perkin Elmer (Buckinghamshire, UK). Rat insulin standard was from  
16    Novo Industria, Copenhagen, Denmark. All other reagents including DPP-4, from porcine  
17    kidney ( $\geq 10$  units/mg protein), Alcalase® 2.4 L and Flavourzyme® 500 L supplied by Sigma  
18    Chemical Company Ltd. (Wicklow, Ireland). Air-dried milled (5 mm) *P. palmata* sample was  
19    purchased from Irish Seaweeds Ltd., Belfast, Co. Antrim, N. Ireland. The macroalgae was  
20    further milled with a Cyclotec™ Mill (1 mm screen, FOSS Tecator AB, Hoganas, Sweden)  
21    and stored at RT.

22

### 23    **Preparation of crude aqueous soluble protein extracts**

Crude aqueous and alkaline soluble protein extracts were prepared using the method described previously [32]. Milled *P. palmata* powder was suspended at a mass:volume ratio of 1:20 ((w/v), 1 kg:20.0 l) and gently agitated at room temperature for 3 h. The supernatant containing the aqueous soluble protein was acquired following centrifugation at 4,190 x g (Sorvall RC6 Plus, Fisher Scientific, Dublin, Ireland) for 15 min at RT. The pellet was resuspended in 0.12 M NaOH (1:15 (w/v)) and gently agitated for 1 h at RT and supernatant containing the alkaline soluble protein was acquired following centrifugation. The pellet was subjected to a second alkaline extraction, and both supernatants combined. A double isoelectric precipitation step was utilised to semi-purified and concentrated aqueous (pH 2.5) and alkaline (pH 4.0) soluble protein components. The precipitated protein pellets obtained following the second isoelectric precipitation were resuspended in dH<sub>2</sub>O to a protein concentration of ~ 2.4% (w/v) and combined. Protein concentration was determined by the modified Lowry protein quantification method as described previously [35]. Samples were analysed in triplicate.

## Enzymatic hydrolysis of macroalgal proteins and simulated gastrointestinal digestion

Macroalgal protein was hydrolysed as described previously [32]. A 2% (w/v) protein solution was preheated to 50°C and adjusted to pH 7.0 and hydrolysed with Alcalase 2.4 L, Alcalase 2.4 L and Flavourzyme 500 L, bromelain and Promod 144 MG at an enzyme:substrate (E:S) ratio of 1:100 (w/w or v/w) for 4 h at 50°C. The reaction was maintained at pH 7.0 using a pH-stat (842 Titrando, Metrohm, Switzerland) and enzyme inactivated by heating at 90°C for 20 min. A control protein sample, containing no proteolytic enzyme, was treated in the same manner. All samples were subsequently freeze-dried (FreeZone 18L, Labconco, MO, USA) and stored at -20°C.

To assess oral bioavailability, PPPH's were subjected to simulated gastrointestinal digestion (SGID), described previously [36]. In brief, unhydrolysed protein controls and hydrolysates were diluted to 2.0% (w/v) protein in water and incubated at 37°C and pH 2 for 90 min with pepsin at an E:S of 1:40 (w/w). The samples were adjusted to pH 7 and subjected to heat inactivation at 90°C for 20 min. The samples were incubated for a further 150 min at 37°C with Corolase PP (E:S of 1% (w/w)). SGID samples were heat inactivated and all samples were subsequently freeze-dried and stored at -20°C.

## Physicochemical characterisation of PPPH

The molecular mass distribution profile of the hydrolysates and their SGID samples were determined by gel permeation-high performance liquid chromatography (GP-HPLC) as described previously [37]. The amino nitrogen content of PPPH was estimated by the TNBS method with absorbance readings taken at 350 nm [38]. Samples were analysed in triplicate.

## Insulin secretion studies in clonal pancreatic cells

Insulinotropic effects of PPPH and SGID samples were measured *in vitro* using clonal pancreatic BRIN-BD11 cells [39]. BRIN-BD11 cells ( $1.5 \times 10^5$  cells/well) were incubated for 20 min with a range of PPPH concentrations (0.039–2.5 mg/ml) in the presence of 5.6 mM glucose at 37°C. Following incubation, supernatant (900 µL) was withdrawn and frozen at -20°C until required. Insulin was quantified using a dextran-coated charcoal radioimmunoassay (RIA), using crystalline rat insulin standard, guinea-pig anti-porcine antiserum (1:30,000 dilution) and  $^{125}\text{I}$ -bovine standard (10,000 cpm), described previously [40]. The concentration of insulin in each sample was determined in duplicate from the prepared insulin standard curve ranging from 0.039-20 ng/ml.

## Cellular toxicity via MTT assay

To determine cytotoxicity of PPPH and SGID samples on BRIN-BD11 cells, the MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide) assay was employed. A fixed dose of PPPH was prepared in Krebs Ringer bicarbonate buffer (KRBB) buffer supplemented with 5.6 or 16.7 mM glucose. Upon completion of co-incubation, KRBB was removed and cells washed with HBSS. Growth media (100 µl) was added to each well of a 96 well plate and further supplemented with 20 µl of MTT solution (5 mg/ml stock). Plates were incubated for 2 h in a modified atmosphere (95% O<sub>2</sub>, 5% CO<sub>2</sub>) tissue culture incubator at 37°C. MTT/growth media was aspirated and washed for a final time with HBSS. The formazan crystals developed were then dissolved using 100 µl of DMSO and the plate agitated at RT for 5 min. Plates were read on a spectrophotometer with absorbance set at 570 nm.

## Quantification of DPP-4 inhibition

DPP-4 inhibition was determined as described previously [33]. Activity was expressed as IC<sub>50</sub> values for three independent replicates (n=3). Diprotin A was used as a positive control.

## *In vitro* GLP-1 secretion from GLUTag cells and GIP secretion from STC-1 cells

*In vitro* effects of the PPPH and the SGID samples on GLP-1 secretion were measured using the murine enteroendocrine GLUTag cell line [41], kindly gifted by Prof. Fiona Gribble/Daniel Drucker. Cells were cultured in high glucose (25 mM) Dulbecco's Modified Eagle's Medium (without glutamine), as described previously [42]. Cells were seeded into 24 well plates (1.5 x 10<sup>5</sup> cells/well) attaching over 36 h at 37°C. Following a pre-incubation step (1.1 mM glucose solution in KRBB for 40 min at 37°C), cells were incubated with the PPPH and SGID samples

(2.5 mg/ml) prepared in 2 mM glucose followed by 2 h incubation at 37°C. Thereafter, 800 µl of supernatant was collected and subsequently used to measure total GLP-1 release by ELISA (Millipore, Hertfordshire, UK) as per manufacturer's protocol.

STC-1 cells differentiate by secreting satiety and glucose homeostatic hormones such as CCK, GIP, PYY, GLP-1 and GLP-2 [43]. The experimental procedure was similar to the GLUTag screening procedure. After 2 h co-incubation, 800 µl of the supernatant was aspirated and stored at -20°C before quantification using a GIP ELISA (Millipore, Hertfordshire, UK) as per manufacturer's protocol.

## Glucose uptake study using differentiated adipocytes

Adipocyte (3T3-L1) cells were obtained from the American Type Culture Collection (ATCC, Manassas, Virginia, USA). 3T3-L1 cells were seeded in a 96, black-walled, clear bottom plates (2 x 10<sup>4</sup> cell/well). Cells were maintained with Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% (v/v) heat inactivated FBS. Cells were incubated for a further 2 days and then differentiated in DMEM containing 10% FBS, 15 µg/mL insulin, 1 µM dexamethasone and 0.5 mM 3-isobutyl-1-methylxanthine (IBMX). Cells were cultured in DMEM containing 10% FBS and 15 µg/mL of insulin. Cells were treated with the test sample (100 µl) or control which were supplemented in glucose-free culture medium containing 150 µg/ml fluorescently tagged 2-deoxyglucose analogue (2-NBDG) and incubated for 20 min. Plates were centrifuged for 5 min at 400 x G at RT. Supernatant was aspirated, and cells washed with 200 µl cell-based assay buffer followed by further centrifugation for 5 min. Wash buffer was removed and 100 µl of cell-based assay buffer was added to all wells and the fluorescence was read immediately at 485 nm with emission measured at 535 nm using the FlexStation scanning fluorimeter (Molecular Devices, Sunnyvale, CA, USA).

## Mechanistic studies using BRIN-BD11 cells

Effect of PPPH's (2.5 mg/ml) on changes in membrane potential and intracellular calcium concentration  $[Ca^{2+}]$  were determined fluorometrically utilising monolayers of BRIN-BD11 cells and Flex membrane potential and calcium assay kits (Molecular Devices, Sunnyvale, CA, USA), as previously described [44]. Assay choice was based on the knowledge that increased intracellular  $[Ca^{2+}]$  is the primary insulin secretory signal, while cAMP signalling-dependent mechanisms are also critical for incretin-mediated insulin release [72]. Control cultures were 30 mM KCl and 10 mM alanine in the presence of 5.6 mM glucose. Fluorometric data was acquired using a FlexStation scanning fluorimeter utilizing an integrated fluid transfer workstation (Molecular Devices, Sunnyvale, CA, USA). The effect of the PPPH's subjected to SGID on the production of cAMP was also assessed in BRIN-BD11 cells. Cells were seeded ( $1.5 \times 10^5$  cells/well) into 24-well plates and incubated overnight. Cells were washed with HBSS before incubation (20 min, 37°C) with the PPPH (2.5 mg/ml) in the presence of 200  $\mu$ M IBMX. Culture media was removed, cells lysed and the cAMP concentration in lysates was determined using a cAMP detection kit (R&D Systems Parameter, Abingdon, UK).

## Acute *in vivo* effects of a PPPH on glucose tolerance and satiety

NIH Swiss mice (Harlan UK Ltd., Blackthorne, UK) were employed for acute *in vivo* experiments. Animals (10-12 week old) were maintained in an environmentally controlled laboratory at  $22 \pm 2^\circ\text{C}$  with a 12 h dark and light cycle with *ad libitum* access to standard rodent diet (10% fat, 30% protein and 60% carbohydrate; Trouw Nutrition, Northwich, UK) and drinking water. Acute glucose-lowering and insulin releasing properties of Alcalase/Flavourzyme PPPH was determined in age-matched groups (n=8) of overnight fasted mice, who received an oral gavage of either glucose alone (18.8 mmol/kg body

combination with PPPH (100 mg/kg bw). Blood glucose was measured using an Ascencia Contour blood glucose meter (Bayer Healthcare, Newbury, UK) and samples were collected via tail vein bleed in chilled fluoride/heparin micro-centrifuge tubes (Sarstedt, Numbrecht, Germany) and centrifuged at 13,000 rpm for 10 min. Plasma was aliquoted and stored at -20°C until required for insulin determination using a modified dextran-coated charcoal RIA [40].

Satiating effect of Alcalase/Flavourzyme PPPH was assessed in male HsD:Ola TO mice (10-12 weeks, Envigo, Blackthorn, UK), maintained as above. Animals had *ad libitum* access to food for 1 week. This was reduced to 10 h of food availability daily on week 2, with further reduction to 6 h daily by week 3. Finally, on week 4 and for the duration of the satiety studies, food availability was strictly maintained at 3 h daily (10.00-13.00 h). Animals (n=8) received an oral dose of saline (0.9 % NaCl) alone, or in combination with PPPH (100 mg/kg bw) immediately prior to regular food access at 10.00 h. Food intake was measured at 30 min intervals up to 180 min.

## Statistical analysis

Results were analysed using GraphPad PRISM 5.0 (San Diego, CA, USA), with data presented as mean  $\pm$  SEM. Comparative analyses between groups were carried out using Student's unpaired *t*-test, one-way ANOVA with a Bonferroni *post hoc* test, or a two-way repeated measures ANOVA with a Bonferroni *post hoc* test where appropriate. Results were deemed significant once  $p < 0.05$ .

## Results

### Insulin secretion and cell viability following PPPH co-incubation with BRIN-BD11 cells

Insulin secretion was determined over a 20 min co-incubation with PPPH supplemented glucose. Baseline insulin secretion was established utilising KRBB buffer supplemented with basal 5.6 mM or elevated 16.7 mM glucose. Several PPPH's, subjected to different hydrolysis conditions, were employed along with SGID equivalents over an identical concentration range (0.0195–2.5 mg/ml). Aqueous/alkaline protein isolate which was subjected to similar hydrolysis condition to that of PPPH's, albeit without the addition of enzyme, was employed as a control and elicited elevated ( $p<0.01$  -  $p<0.001$ ) insulin secretion at 1.25 and 2.5 mg/ml at basal glucose concentration (Fig. 1A). Interestingly, its SGID equivalent presented with surprisingly high insulintropic activity ( $p<0.01$ - $p<0.001$  at  $> 0.312$  mg/ml) when tested at 16.7 mM glucose concentration in BRIN-BD11 cells (Fig. 1B). While potentially anomalous, improved potency ( $p<0.01$ - $p<0.001$  at  $> 0.156$  mg/ml) following SGID highlights the importance of more complete hydrolysis to the insulintropic effect.

The inverse was true for PPPH's. Alcalase PPPH stimulated insulin secretion ( $p<0.001$ ) from 0.312 mg/ml and above (Fig. 1C). However, following SGID, the effect on insulintropic potency was negatively impacted, with bioactivity ( $p<0.05$  to  $p<0.01$ ) observed from 1.25 mg/ml and above (Fig. 1D). Alteration of the hydrolysis medium for Alcalase/Flavourzyme PPPH improved potency, with augmented ( $p<0.01$  to  $p<0.001$ ) insulin secretion from 0.312 mg/ml and above (Fig. 1E). Post-SGID, Alcalase/Flavourzyme PPPH displayed improved potency, increasing ( $p<0.001$ ) insulin secretion at 0.625 mg/ml or above (Fig. 1F). Of other digestion conditions, the bromelain PPPH exhibited promising insulin secretory actions, with improvements ( $p<0.05$  -  $p<0.001$ ) compared to baseline at 0.078 mg/ml or above (Fig. 1G).



SGID negatively impacted efficacy; however, potency was still impressive, enhancing ( $p<0.001$ ) insulin secretion from 1.25 mg/ml or higher (Fig. 1H). Somewhat unexpectedly, the Promod PPPH (Fig. 1I) and its SGID equivalent (Fig. 1J) displayed bioactivity over an identical concentration range (0.625 - 2.5 mg/ml) with only the magnitude of the increased insulin secretion being slightly impacted following SGID.

PPPH's were further tested in the presence of 16.7 mM glucose. While potency was slightly altered, the magnitudes and trends involving SGID were largely the same for the aqueous, Alcalase, Alcalase/Flavourzyme PPPH's and their SGID counterparts (Fig. 2A-F). The potency of both bromelain and Promod PPPH's at elevated glucose was reduced, with respect to the 5.6 mM glucose data, but they retained a dose-dependent effect above 0.625 (Fig. 2G,I). SGID enhanced the potency of these PPPH's in both cases (Fig. 2H,J). Importantly, when tested at the highest concentration, cell viability was not negatively impacted by the inclusion of any PPPH at either 5.6 mM (Fig. 1K) or 16.7 mM glucose (Fig. 2K).

## Preservation of the incretin effect

As shown in Table 1, DPP-4 inhibition significantly increased following digestion with all proteolytic enzymes employed. Greatest inhibition was observed with PPPH's generated with Alcalase/Flavourzyme or Alcalase alone, achieving DPP-4  $IC_{50}$  values of  $0.70 \pm 0.02$  and  $0.94 \pm 0.10$  mg/ml, respectively. Bromelain and Promod PPPH's had lower DPP-4 inhibitory activity with  $IC_{50}$  values of  $1.34 \pm 0.05$  and  $1.23 \pm 0.05$ , respectively.  $IC_{50}$  values of the control and PPPH's were significantly altered by SGID. The DPP-4 inhibitory activity mediated by the control and bromelain and Promod PPPH's increased ( $p<0.05$ ) following SGID. In contrast, the DPP-4 inhibitory activity with Alcalase/Flavourzyme and Alcalase PPPH's decreased

( $p<0.05$ ) following SGID. Thus, in the latter case peptides eliciting high DPP-4 inhibitory activity in the hydrolysate were degraded during SGID.

## Promotion of the incretin effect

The effects of PPPH upon GLP-1 and GIP secretion was investigated via acute exposure of enteroendocrine GLUTag and STC-1 cell lines, respectively. Positive controls, glutamine (10 mM), forskolin (10 mM) and GIP ( $10^{-6}$  M) returned from 2- to 4-fold ( $p<0.05$  -  $p<0.001$ ) increases in GLP-1 secretion when compared to basal glucose (2 mM) control (Fig. 3A). Likewise, palmitic acid (500  $\mu$ M) and glutamine (10 mM) showed a 4-fold increase ( $p<0.001$ ) in GIP secretion compared to glucose control (Fig. 3B). PPPH's (2.5 mg/ml) were subsequently co-incubated with 2 mM glucose to investigate their effects on hormone secretion.

The aqueous/alkaline protein control elicited a 3-fold increase ( $p<0.001$ ) in GLP-1 secretion compared to basal glucose control (Fig. 3A). Following SGID, bioactivity was reduced ( $p<0.05$ ); however, GLP-1 secretion remained elevated ( $p<0.01$ ) versus the 2 mM glucose control (Fig. 3A). Interestingly, the inverse was true for GIP secretion, where the protein control elicited a 4-fold ( $p<0.001$ ) increase, but the SGID equivalent led to 5.8-fold ( $p<0.001$ ) increase in GIP secretion compared to glucose control (Fig. 3B). Alcalase PPPH displayed a 2-fold ( $p<0.001$ ) increase in GLP-1 secretion, but post-SGID it failed to raise secretion beyond the glucose control (Fig. 3A). For GIP secretion, Alcalase PPPH resulted in a 6-fold ( $p<0.001$ ) increase which was improved post SGID, with an 8-fold ( $p<0.001$ ) upregulation (Fig. 3B). Little change was observed following addition of Flavourzyme, with the hydrolysate promoting a 2.2-fold increase in GLP-1 secretion ( $p<0.001$ ) accompanied by a loss of bioactivity following SGID (Fig. 3A). Unexpectedly, GIP secretion was impacted by

Alcalase/Flavourzyme digestion, whereby the secretory activity was mildly elevated (1.2-fold;  $p<0.01$ ); however, following SGID, a significant reduction ( $p<0.01$ ) in secretion was observed (Fig. 3B). Bromelain PPPH produced 2-fold ( $p<0.001$ ) rise in GLP-1 secretion but only retained a 1.4-fold ( $p<0.05$ ) secretory response post SGID (Fig. 3A). Bromelain PPPH and its SGID counterpart increased GIP secretion 5-fold ( $p<0.001$ ) and 3.5-fold ( $p<0.001$ ), respectively, retaining relatively high bioactivity post SGID (Fig. 3B). The same was largely true for Promod PPPH, displaying a 1.6-fold ( $p<0.001$ ) increase in GLP-1 secretion which was reduced to 1.4-fold ( $p<0.05$ ) following SGID (Fig. 3A). Again, GIP secretion was relatively well-retained with 6-fold ( $p<0.001$ ) and 3.8-fold ( $p<0.001$ ) improvements in hormone secretion beyond control culture for Promod PPPH and its SGID equivalent, respectively (Fig. 3B).

### Mechanistic consequences of co-incubation with PPPH

The cellular consequences following co-incubation of 2.5 mg/ml of PPPH were investigated in BRIN-BD11 cells supplemented with 5.6 mM glucose. Specifically, cyclic adenosine monophosphate (cAMP), intracellular calcium ( $[Ca^{2+}]_i$ ) and membrane potential were tested under identical conditions. cAMP was initially used to screen PPPH's. The positive controls, 16.7 mM glucose and GLP-1 ( $10^{-6}$  M), elevated ( $p<0.05$  and  $p<0.001$ , respectively) intracellular cAMP versus 5.6 mM glucose control (Fig. 4A). All six non-SGID, PPPH's raised cAMP production ( $p<0.05$  -  $p<0.001$ ) in BRIN-BD11 cells (Fig. 4A), while only SGID, Alcalase and SGID, Promod PPPH's failed to stimulate cAMP above the 5.6 mM control (Fig. 4A). Notably, Alcalase/Flavourzyme PPPH and its SGID counterpart, elicited the greatest rise in intracellular cAMP, upregulating 1.7-fold ( $p<0.001$ ) compared to 5.6 mM glucose control (Fig. 4A). As a result of constraints over assay availability, combined with the limited

availability of SGID sample, only Alcalase/Flavourzyme PPPH was investigated further with respect to cellular signalling.

In terms of intracellular  $\text{Ca}^{2+}$  mobilisation, the positive control, alanine (10 mM), elicited a 15-fold increase ( $p<0.001$ ) in calcium mobilisation compared to the 5.6 mM glucose control culture (Fig. 4B,C). Alcalase/Flavourzyme PPPH greatly surpassed ( $p<0.001$ ) the positive control, increasing  $\text{Ca}^{2+}$  mobilisation 80-fold ( $p<0.001$ ) versus the 5.6 mM glucose control (Fig. 4B,C). With respect to membrane potential, potassium chloride (KCl, 30 mM) a potent membrane potentiating insulinotropic electrolyte, caused a 75-fold increase ( $p<0.001$ ) in membrane potential during acute co-incubation with 5.6 mM glucose (Fig. 4D,E). Further to increased  $\text{Ca}^{2+}$ , Alcalase/Flavourzyme PPPH returned a 125-fold peak increase ( $p<0.001$ ) in membrane potential versus 5.6 mM glucose control in BRIN-BD11 cells (Fig. 4D,E).

### Effects of PPPH on *in vitro* glucose uptake

The 3T3-L1 cell line was investigated following trans-differentiation from fibroblast to adipocyte cells. Apigenin control culture evoked a significant reduction ( $p<0.001$ ) of glucose uptake, via the inhibition of the GLUT-1 receptor, with the inverse for low (1 nM) and high (100 nM) insulin, causing 1.4- and 1.8-fold increases ( $p<0.001$ ) in glucose uptake, respectively (Fig. 5A). All PPPH's and their SGID equivalents, were employed either alone at 2.5 mg/ml or in combination with basal insulin (1 nM).

Interestingly, while the protein control failed to show a significant increase in glucose on its own, in the presence of insulin a 1.5-fold increase ( $p<0.05$ ) was observed (Fig. 5B). The inverse was true following SGID causing significantly increased ( $p<0.05$ ) glucose uptake on its own, with surprising loss of effect in the presence of insulin (Fig. 5B). When co-incubated alone, Alcalase PPPH and its SGID equivalent both caused comparable, 1.5-fold ( $p<0.001$ ),

increases in glucose uptake; however, when co-incubated with insulin, no additive effect was displayed (Fig. 5C). Alcalase/Flavourzyme PPPH, when incubated alone, caused a 1.8-fold increase ( $p<0.001$ ) in glucose uptake, with co-incubation with insulin not impacting greatly, demonstrating a 1.7-fold increase ( $p<0.001$ ) in uptake (Fig. 5D). Similarly, post-SGID, elicited a 1.38-fold ( $p<0.05$ ) rise in glucose uptake both with and without insulin present (Fig. 5D). Bromelain PPPH failed to improve glucose uptake when incubated on its own, with notable improvement presented as a 1.5-fold increase ( $p<0.01$ ) following insulin co-incubation (Fig. 5E). However, following SGID, the sample failed to produce an improvement in glucose uptake in the absence or presence of insulin (Fig. 5E). Similarly, Promod PPPH was only effective in the presence of insulin, when glucose uptake increased by 1.5-fold ( $p<0.05$ ), but this was lost following SGID, with the sample having no effect either alone or in the presence of insulin (Fig. 5F).

#### Acute *in vivo* effects following oral administration of PPPH

Following initial dose-response investigations (see Supplementary Figures), a 100 mg/kg/bw dose of Alcalase/Flavourzyme PPPH was employed for *in vivo* investigations. When co-administered with 18.8 mM glucose, PPPH significantly curbed rises in blood glucose at 60 min ( $p<0.01$ ), as well as 90 and 120 min ( $p<0.001$ ), when compared to glucose-only control (Fig. 6A). Additionally, the area under the curve (AUC) was significantly reduced ( $p<0.01$ ) compared to control (Fig. 6B). Positive glycaemic effects appear to be relatively short-lasting, with a delayed oral glucose challenge showing no anti-hyperglycaemic efficacy when delivered 4, 8 or 12 h post PPPH administration (data not shown).

1        Additionally, when co-administered with saline, PPPH (100 mg/kg) demonstrated  
2        significant reductions ( $p<0.05$  to  $p<0.001$ ) in food intake (16-20% from 90 to 180 min) when  
3        tested in diet-restricted animals (Fig. 6C).

## 5        **Discussion**

6        While accepted that an increase in dietary protein intake can have positive effects on glycaemic  
7        control for T2DM patients [10]; expanding population sizes, coupled with pre-existing social  
8        inequality, continue to drive global protein malnutrition [45-47]. It is predicted that the global  
9        human population will reach 9 billion by 2050, meaning trends in population expansion are  
10       unlikely to be curbed. Thus, greater emphasis must be placed on ensuring food security [48].  
11       Novel sources are being pursued to address this, one such being the exploration of marine  
12       sources like seaweeds, industry off-cuts and underutilised, or low value fish species, many of  
13       which are currently used as animal and farmed fish/shellfish feed [31,49,50]. Algal proteins  
14       are of particular interest due to a favourable amino acid composition, containing all nine  
15       essential amino acids [51], combined with a relatively high protein content, particularly for red  
16       seaweeds which often contain  $\geq 30\%$  of protein by dry weight [52].

17       It was recently established that chronic administration of an Alcalase/Flavourzyme PPPH  
18       can improve glycaemic control in STZ-induced diabetic mice [34]. Therefore, utilising an array  
19       of *in vitro* and acute *in vivo* techniques, the present study aimed to uncover causative  
20       mechanisms of PPPH-induced benefits in diabetes, investigating Alcalase/Flavourzyme PPPH  
21       while taking the opportunity to compare it to a number of other enzymatically produced  
22       PPPH's.

23       Initial screening involved assessment of insulinotropic activity of PPPH's in BRIN-  
24       BD11 cells, which have been previously utilised in the screening of enzymatically-produced

1 protein hydrolysates [25-27]. We hypothesised that activity would rely on the hydrolysis  
2 process, specifically the duration of hydrolysis and the protease used, ultimately leading to a  
3 different range of bioactive peptides within crude mixtures [53]. Our data appears to support  
4 this, particularly with respect to the choice of enzyme and the extent to which it hydrolyses *P.*  
5 *palmata* proteins and the MW of peptides therein. As reflected in the molecular distribution  
6 profiles (Supplementary Table 1), and accompanying GP-HPLC profiles (Supplementary Fig.  
7 4), both Alcalase and Alcalase/Flavourzyme mediated a higher extent of hydrolysis of *P.*  
8 *palmata* proteins than bromelain and Promod. Also, all hydrolysates were further hydrolysed  
9 during the *in vitro* digestion process, which had either a positive or negative impact on  
10 bioactivity.

11       Insulinotropic effects of the protein control, were vastly improved following SGID, while  
12 the insulinotropic potency of Alcalase and Alcalase/Flavourzyme PPPH's subjected to SGID  
13 was reduced rather than totally abolished, an encouraging finding. Intriguingly, particularly at  
14 16.7 mM glucose, bromelain and Promod PPPH's showed unaltered or even improved  
15 bioactivity post SGID. Further analysis may uncover the possibility that glycation of specific  
16 components can affect their bioactivity, with the data suggesting that, for most PPPH's, the  
17 potency of insulinotropic effect was greater at 16.7 mM than at 5.6 mM glucose. Given  
18 insulinotropic peptides, such as GIP, are demonstrated to have improved bioactivity following  
19 N-terminal glycation [54,55], this may be plausible, but remains a hypothesis. Furthermore,  
20 the favourable MTT data showing no cellular cytotoxicity, demonstrated that the increased  
21 insulin release induced by PPPH was not a result of beta-cell lysis [56].

22       It is well established that both crude hydrolysates [32], and isolated peptide sequences  
23 [33], derived from *P. palmata*, have excellent DPP-4 inhibitory properties. As such, for the  
24 present set of PPPH's, these findings have been corroborated. Furthermore, the effects were  
25 largely retained or even improved upon following SGID, indicating extensive oral

1 bioavailability [25,27]. While likely a less important factor in *in vitro* insulin secretion than in  
2 *in vivo* scenarios, previously identified presence of intra-islet DPP-4 [73], as well as  
3 expression within beta-cells themselves [74], means the influence of DPP-4 inhibitory  
4 effects to the insulintropic effects of PPPH's cannot be completely ruled out, but certainly  
5 warrant further investigation. The clinical success of DPP-4 inhibition is mediated through  
6 preservation of the incretin effect, and this mechanism is well established in management of  
7 T2DM [23,57-59]. The present data demonstrates that: beyond preservation of the incretin-  
8 effect, crude PPPH's can directly influence incretin-mediated glycaemic improvements via  
9 the stimulation of GLP-1 and GIP secretion [60]. Thus, all PPPH's enhanced both *in vitro*  
10 GLP-1 and GIP secretion in their unaltered states, a finding supported by previous findings of  
11 marine protein-hydrolysate-induced GLP-1 secretion [25,27]. Intriguingly, the effects on GIP  
12 secretion were more resistant to SGID than those of GLP-1, with the effect at least retained  
13 for every hydrolysate except Alcalase/Flavourzyme PPPH. It is important to note that,  
14 given the crude nature of the hydrolysate mixtures, differing small MW peptides or free  
15 amino acid components may be responsible for each effect.

16 *In vitro* analysis of mechanistic consequences of PPPH co-incubation further highlight  
17 the ability of PPPH to influence insulin release. Membrane depolarisation, calcium  
18 mobilisation and generation of cAMP were employed to provide a general overview of activity.  
19 As such, cAMP upregulation suggests that small MW peptides may be stimulating secretion of  
20 hormones such as insulin, GLP-1 and GIP via G-coupled protein receptor activation [61]. These  
21 receptors are only activated by extracellular stimuli and can have potent cellular activation via  
22 promotion of internal signalling cascades [62,63]. For example, activation of the GLP-1r via  
23 GLP-1 promotes a significant increase in cAMP production and glucose dependent insulin  
24 secretion [63]. Indeed, each PPPH elicited cAMP elevation beyond baseline, with varying  
25 degrees of effect. In particular, Alcalase/Flavourzyme PPPH exhibited the most potent



elevation, with full retention following SGID. Furthermore, this hydrolysate greatly influenced both membrane depolarisation and calcium mobilisation, beyond positive controls. While the direct effects on intracellular signalling appear to be multifaceted [62-64], further investigation is required to identify whether the bioactivity emanates from a potent singular peptide entity within the crude mixture, or if it arises from synergistic mechanisms of various peptides or amino acids [65].

3T3-L1 adipocyte cells were utilised to examine the ability of the PPPH and SGID samples to stimulate glucose uptake. These cells exhibit all the components of insulin receptor and signal transduction cascade and are frequently used to investigate insulin mediated glucose transport [66]. It is believed that compensatory upregulation of insulin production and secretion can result in impaired glucose transport into liver, skeletal muscle, and adipose tissue in T2DM [67]. Furthermore, it is postulated that dysfunctional peripheral glucose uptake, may contribute to insulin resistance in skeletal muscle [68]. The present data unveils a positive role for PPPH in glucose transport, with Alcalase and Alcalase/Flavourzyme PPPH's increasing glucose uptake, independent of insulin, with the effect partially surviving SGID. Such findings may be related to the acute glucose tolerance test data. Thus, oral administration of Alcalase/Flavourzyme PPPH improved the oral glucose tolerance significantly from 60 min onwards of challenge initiation. While thought to be multifactorial in nature, improved glucose uptake could be a contributing factor [69]. Insulin determination was not possible during these experiments; however, previous findings have demonstrated that chronic administration of an equivalent dose of this hydrolysate (split into twice daily 50 mg/kg/bw dosage) improved non-fasting insulin levels, with accompanying improvements in circulating glucose and HbA<sub>1c</sub>, over the treatment period [34]. Additionally, we highlight a role for PPPH administration in satiety, with notable reduction from baseline evident from 90-180 min inclusive when tested in food deprived trained mice, again supported by previous findings [34].

1        Taken together, this dataset fortifies interest in PPPH with regards to management of  
2        T2DM. Knowledge of the previously established roles in DPP-4 inhibition [32] and partial  
3        reversal of STZ-induced diabetes [34] have been expanded upon. Here, we show clear roles for  
4        PPPH in direct insulin and incretin secretion, supported by cellular signalling data, as well as  
5        improvements in glucose utilisation and tolerance now identified. Given it is now accepted  
6        that particular combinations of individual amino acids, such as leucine, alanine and  
7        glutamine, at supraphysiological concentrations can augment insulin secretion [72], we  
8        postulate that such a mechanism may be at play here with crude hydrolysate mixtures, thus  
9        future work may isolate, identify and characterise specific peptides, single amino acids or  
10       combinations thereof, small molecules or lipids from the most consistently  
11       promising of the hydrolysates, Alcalase/Flavourzyme PPPH [33]. This hydrolysate was  
12       particularly promising given its direct insulintropic effects, accompanied by promising  
13       effects on the intracellular mechanisms linked to insulin release, along with positive effects  
14       on glucose uptake.

15       The study possesses some limitations, primarily that limited SGID sample availability  
16       did not permit inclusion of all test hydrolysates in our mechanistic investigations. In the  
17       present preliminary study, the findings observed herein on glycaemic control and insulin  
18       secretion could arise due to multiple different actions mediated by the PPPH peptides.  
19       These might involve for example, the inhibition of DPP-4 prolonging the bioactivity of  
20       GLP-1 and GIP incretin hormones or insulintropic actions of free amino acids. However,  
21       extensive detailed mechanistic studies would be required to elucidate the various  
22       mechanisms, which is beyond the scope of the current study. Furthermore, a chronic study  
23       with isolated peptides would be warranted to increase our knowledge on the bioactivity  
24       residing in PPPH. Additionally, given all the hydrolysates assessed herein inhibit  
angiotensin converting enzyme (ACE) (data not shown) combined with our clearly  
identified effects on satiety, it may be interesting to

investigate the influence PPPH can play on regulation of satiating hormones like  
cholecystokinin (CCK) and PYY(3-36), both of which are substrates for ACE [70,71].

## **Conflict of interest**

On behalf of all authors, the corresponding author states that there is no conflict of interest.

## **Ethics statement**

All animal protocols were approved by the Ulster University, Animal Welfare and Ethical  
Review (AWERB) Committee and performed according to the UK Animals (Scientific  
Procedures) Act 1986 and EU Directive 2010/63EU.

## **Authors' contributions**

FOH and PHR designed the experiments, CMcL, VP and PHR performed experiments and  
collected and analyzed data, CMcL and RL wrote the draft manuscript. PA, EMcS, SS, RF and  
FOH supervised the work and read and edited the manuscript.

## References

1. Wilcox G (2005) Insulin and insulin resistance. Clin. Biochem. Rev 26: 19-39.
2. Donath MY, Ehses JA, Maedler K, et al (2005) Mechanisms of  $\beta$  -cell death in type 2 diabetes. Diabetes 54:108-113. [https://doi.org/10.2337/diabetes.54.suppl\\_2.S108](https://doi.org/10.2337/diabetes.54.suppl_2.S108)
3. Lund A, Bagger JJ, Albrechtsen NJW, et al (2016) Evidence of extrapancreatic glucagon secretion in man. Diabetes 65:585–597. <https://doi.org/10.2337/db15-1541>
4. International Diabetes Federation *IDF Diabetes Atlas, 9th edn*. Brussels, Belgium: 2019. Available at: <https://www.diabetesatlas.org>
5. Cannon A, Handelsman Y, Heile M, Shannon M (2018) Burden of Illness in Type 2 diabetes mellitus. J. Manag. Care Spec. Pharm. 24:S5–S13. <https://doi.org/10.18553/jmcp.2018.24.9-a.s5>
6. Muggeo M (1998) Accelerated complications in Type 2 diabetes mellitus: the need for greater awareness and earlier detection. Diabet. Med. 15 Suppl 4:S60-62. [https://doi.org/10.1002/\(sici\)1096-9136\(1998120\)15:4+<s60::aid-dia736>3.3.co;2-a](https://doi.org/10.1002/(sici)1096-9136(1998120)15:4+<s60::aid-dia736>3.3.co;2-a)
7. Tuomilehto J, Schwarz P, Lindström J (2011) Long-term benefits from lifestyle interventions for type 2 diabetes prevention: Time to expand the efforts. Diabetes Care 34:S210–S214. <https://doi.org/10.2337/dc11-s222>
8. Forouhi NG, Misra A, Mohan V, et al (2018) Dietary and nutritional approaches for prevention and management of type 2 diabetes. BMJ 361:k2234. <https://doi.org/10.1136/bmj.k2234>
9. Wu G (2016) Dietary protein intake and human health. Food Funct. 7:1251–1265. <https://doi.org/10.1039/c5fo01530h>

10. Gannon MC, Nuttall FQ, Saeed A, et al (2003) An increase in dietary protein improves the blood glucose response in persons with type 2 diabetes. *Am. J. Clin. Nutr.* 78:734–741. <https://doi.org/10.1093/ajcn/78.4.734>
11. Zhao W, Luo Y, Zhang Y, et al (2018) High protein diet is of benefit for patients with type 2 diabetes: An updated meta-analysis. *Medicine* 97:e13754. <https://doi.org/10.1097/MD.00000000000013149>
12. Newsholme P, Krause M (2012) Nutritional regulation of insulin secretion: Implications for diabetes. *Clin. Biochem. Rev.* 33:35–47 <https://doi.org/01598090>.
13. Oseguera-Toledo ME, González de Mejía E, Reynoso-Camacho R, et al (2014) Proteins and bioactive peptides. *Nutrafoods* 13:147–157. <https://doi.org/10.1007/s13749-014-0052-z>
14. Nongonierma AB, Fitzgerald RJ (2013) Dipeptidyl peptidase IV inhibitory and antioxidative properties of milk protein-derived dipeptides and hydrolysates. *Peptides* 39:157–163. <https://doi.org/10.1016/j.peptides.2012.11.016>
15. Nongonierma AB, FitzGerald RJ (2013) Dipeptidyl peptidase IV inhibitory properties of a whey protein hydrolysate: Influence of fractionation, stability to simulated gastrointestinal digestion and food-drug interaction. *Int. Dairy J* 32:33–39. <https://doi.org/10.1016/j.idairyj.2013.03.005>
16. Nongonierma AB, FitzGerald RJ (2013) Inhibition of dipeptidyl peptidase IV (DPP-IV) by proline containing casein-derived peptides. *J. Funct. Foods* 5:1909–1917. <https://doi.org/10.1016/j.jff.2013.09.012>

17. Nongonierma AB, Fitzgerald RJ (2013) Inhibition of dipeptidyl peptidase IV (DPP-IV) by tryptophan containing dipeptides. Food Funct. 4:1843–1849. <https://doi.org/10.1039/c3fo60262a>
18. Power O, Nongonierma AB, Jakeman P, Fitzgerald RJ (2014) Food protein hydrolysates as a source of dipeptidyl peptidase IV inhibitory peptides for the management of type 2 diabetes. Proc. Nutr Soc 73:34–46. <https://doi.org/10.1017/S0029665113003601>
19. Ahrén B (2007) DPP-4 inhibitors. Best Pract Res Clin Endocrinol Metab 21:517–533. <https://doi.org/10.1016/j.beem.2007.07.005>
20. Holst JJ, Vilsbøll T, Deacon CF (2009) The incretin system and its role in type 2 diabetes mellitus. Mol Cell Endocrinol 297:127–136. <https://doi.org/10.1016/j.mce.2008.08.012>
21. Green B, Gault V, O’Harte F, Flatt P (2005) Structurally Modified Analogues of Glucagon-Like Peptide-1 (GLP-1) and Glucose-Dependent Insulinotropic Polypeptide (GIP) As Future Antidiabetic Agents. Curr Pharm Des 10:3651–3662. <https://doi.org/10.2174/1381612043382774>
22. Gallwitz B (2011) Glucagon-like peptide1 analogues for type 2 diabetes mellitus: Current and emerging agents. Drugs 71:1675–1688. <https://doi.org/10.2165/11592810-000000000-00000>
23. Deacon CF (2019) Physiology and pharmacology of DPP-4 in glucose homeostasis and the treatment of type 2 diabetes. Front Endocrinol 10:80. <https://doi.org/10.3389/fendo.2019.00080>

24. Harnedy PA, Parthasarathy V, McLaughlin CM, et al (2018) Atlantic salmon (*Salmo salar*) co-product-derived protein hydrolysates: A source of antidiabetic peptides. *Food Res Int* 106:598–606. <https://doi.org/10.1016/j.foodres.2018.01.025>
25. Harnedy PA, Parthasarathy V, McLaughlin CM, et al (2018) Blue whiting (*Micromesistius poutassou*) muscle protein hydrolysate with in vitro and in vivo antidiabetic properties. *J Funct Foods* 40:137–145. <https://doi.org/10.1016/j.jff.2017.10.045>
26. Harnedy-Rothwell PA, McLaughlin CM, O’Keeffe MB, et al (2020) Identification and characterisation of peptides from a boarfish (*Capros aper*) protein hydrolysate displaying in vitro dipeptidyl peptidase-IV (DPP-IV) inhibitory and insulinotropic activity. *Food Res Int* 131:108989. <https://doi.org/10.1016/j.foodres.2020.108989>
27. Parthasarathy V, McLaughlin CM, Harnedy PA, et al (2019) Boarfish (*Capros aper*) protein hydrolysate has potent insulinotropic and GLP-1 secretory activity in vitro and acute glucose lowering effects in mice. *Int J Food Sci Technol* 54:271–281. <https://doi.org/10.1111/ijfs.13975>
28. Galland-Irmouli AV, Fleurence J, Lamghari R, et al (1999) Nutritional value of proteins from edible seaweed *Palmaria palmata* (Dulse). *J Nutr Biochem* 10:353–359. [https://doi.org/10.1016/S0955-2863\(99\)00014-5](https://doi.org/10.1016/S0955-2863(99)00014-5)
29. Yuan Y V., Bone DE, Carrington MF (2005) Antioxidant activity of dulse (*Palmaria palmata*) extract evaluated in vitro. *Food Chem* 91:485–494. <https://doi.org/10.1016/j.foodchem.2004.04.039>
30. Yuan Y V., Walsh NA (2006) Antioxidant and antiproliferative activities of extracts from a variety of edible seaweeds. *Food Chem Toxicol* 44:1144–1150. <https://doi.org/10.1016/j.fct.2006.02.002>

31. Harnedy PA, Fitzgerald RJ (2011) Bioactive proteins, peptides, and amino acids from macroalgae. *J. Phycol.* 47:218–232. <https://doi.org/10.1111/j.1529-8817.2011.00969.x>
32. Harnedy PA, FitzGerald RJ (2013) In vitro assessment of the cardioprotective, anti-diabetic and antioxidant potential of *Palmaria palmata* protein hydrolysates. *J Appl Phycol* 25:1793–1803. <https://doi.org/10.1007/s10811-013-0017-4>
33. Harnedy PA, O’Keeffe MB, Fitzgerald RJ (2015) Purification and identification of dipeptidyl peptidase (DPP) IV inhibitory peptides from the macroalga *Palmaria palmata*. *Food Chem* 172:400–406. <https://doi.org/10.1016/j.foodchem.2014.09.083>
34. McLaughlin CM, Sharkey SJ, Harnedy-Rothwell P, et al (2020) Twice daily oral administration of *Palmaria palmata* protein hydrolysate reduces food intake in streptozotocin induced diabetic mice, improving glycaemic control and lipid profiles. *J Funct Foods* 73: 104101. <https://doi.org/10.1016/j.jff.2020.104101>
35. Harnedy PA, FitzGerald RJ (2013) Extraction of protein from the macroalga *Palmaria palmata*. *LWT - Food Sci Technol* 51:375–382. <https://doi.org/10.1016/j.lwt.2012.09.023>
36. Walsh DJ, Bernard H, Murray BA, et al (2004) In vitro generation and stability of the lactokinin  $\beta$ -lactoglobulin fragment (142-148). *J Dairy Sci* 87:3845–3857. [https://doi.org/10.3168/jds.S0022-0302\(04\)73524-9](https://doi.org/10.3168/jds.S0022-0302(04)73524-9)
37. Spellman D, Kenny P, O’Cuinn G, Fitzgerald RJ (2005) Aggregation properties of whey protein hydrolysates generated with *Bacillus licheniformis* proteinase activities. *J Agric Food Chem* 53:1258–1265. <https://doi.org/10.1021/jf048754a>
38. Le Maux S, Nongonierma AB, Barre C, Fitzgerald RJ (2016) Enzymatic generation of whey protein hydrolysates under pH-controlled and non pH-controlled conditions:



Impact on physicochemical and bioactive properties. Food Chem 199:246–251.

<https://doi.org/10.1016/j.foodchem.2015.12.021>

39. McClenaghan NH, Barnett CR, Ah-Sing E, et al (1996) Characterization of a novel glucose-responsive insulin-secreting cell line, BRIN-BD11, produced by electrofusion. Diabetes 45:1132–1140. <https://doi.org/10.2337/diab.45.8.1132>

40. Flatt PR, Bailey CJ (1982) Plasma glucose and insulin responses to glucagon and arginine in Aston ob/ob mice: Evidence for a selective defect in glucose-mediated insulin release. Horm Metab Res 14:127–130. <https://doi.org/10.1055/s-2007-1018945>

41. Drucker DJ, Jin T, Asa SL, et al (1994) Activation of proglucagon gene transcription by protein kinase-A in a novel mouse enteroendocrine cell line. Mol Endocrinol 8:1646–1655. <https://doi.org/10.1210/mend.8.12.7535893>

42. McLaughlin CM, Lampis S, Mechkarska M, et al (2016) Purification, conformational analysis, and properties of a family of tigerinin peptides from skin secretions of the crowned bullfrog *Hoplobatrachus occipitalis*. J Nat Prod 79:2350–2356. <https://doi.org/10.1021/acs.jnatprod.6b00494>

43. Reimann F, Gribble FM (2002) Glucose-sensing in glucagon-like peptide-1-secreting cells. Diabetes 51:2757–2763. <https://doi.org/10.2337/diabetes.51.9.2757>

44. Srinivasan D, Mechkarska M, Abdel-Wahab YHA, et al (2013) Caerulein precursor fragment (CPF) peptides from the skin secretions of *Xenopus laevis* and *Silurana epitropicalis* are potent insulin-releasing agents. Biochimie 95:429–435. <https://doi.org/10.1016/j.biochi.2012.10.026>

45. Müller O, Krawinkel M (2005) Malnutrition and health in developing countries. CMAJ 173:279–286. <https://doi.org/10.1503/cmaj.050342>

46. Semba RD (2016) The rise and fall of protein malnutrition in global health. *Ann. Nutr. Metab.* 69:79–88. <https://doi.org/10.1159/000449175>
47. Henchion M, Hayes M, Mullen A, et al (2017) Future protein supply and demand: Strategies and factors influencing a sustainable equilibrium. *Foods* 6:53. <https://doi.org/10.3390/foods6070053>
48. Cole MB, Augustin MA, Robertson MJ, Manners JM (2018) The science of food security. *npj Sci Food* 2:14. <https://doi.org/10.1038/s41538-018-0021-9>
49. Aneiros A, Garateix A (2004) Bioactive peptides from marine sources: Pharmacological properties and isolation procedures. *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* 803:41–53. <https://doi.org/10.1016/j.jchromb.2003.11.005>
50. Sharkey SJ, Harnedy-Rothwell PA, Allsopp PJ, et al (2020) A narrative review of the anti-hyperglycemic and satiating effects of fish protein hydrolysates and their bioactive peptides. *Mol. Nutr. Food Res.* 64: 2000403. <https://doi.org/10.1002/mnfr.202000403>
51. Bleakley S, Hayes M (2017) Algal proteins: Extraction, application, and challenges concerning production. *Foods* 6:33. <https://doi.org/10.3390/foods6050033>
52. Fleurence J, Levine I (2016) Seaweed in Health and Disease Prevention. Chapter 5. Elsevier Inc. <https://doi.org/10.1016/C2014-0-02206-X>
53. Waseem M, Kumar S, Kumar A (2018) Bioactive peptides. In: Secondary metabolite and functional food components: Role in health and disease. Nova Science Publisher Inc. 6:259–287. <https://doi.org/10.3390/foods6050032>
54. O'Harte FPM, Abdel-Wahab YHA, Conlon JM, Flatt PR (1998) Amino terminal glycation of gastric inhibitory polypeptide enhances its insulinotropic action on clonal

pancreatic B-cells. *Biochim Biophys Acta - Gen Subj* 1425:319–327.

[https://doi.org/10.1016/S0304-4165\(98\)00084-1](https://doi.org/10.1016/S0304-4165(98)00084-1)

55. O'Harte FPM, Mooney MH, Kelly CMN, Flatt PR (2000) Improved glycaemic control in obese diabetic ob/ob mice using n- terminally modified gastric inhibitory polypeptide. *J Endocrinol* 165:639–648. <https://doi.org/10.1677/joe.0.1650639>

56. Janjic D, Wollheim CB (1992) Islet cell metabolism is reflected by the MTT (tetrazolium) colorimetric assay. *Diabetol Clin Exp Diabetes Metab* 35:482–485. <https://doi.org/10.1007/BF02342448>

57. Nauck MA, Heimesaat MM, Orskov C, et al (1993) Preserved incretin activity of glucagon-like peptide 1 [7-36 amide] but not of synthetic human gastric inhibitory polypeptide in patients with type- 2 diabetes mellitus. *J Clin Invest* 91:301–307. <https://doi.org/10.1172/JCI116186>

58. Holst JJ, Deacon CF (2005) Glucagon-like peptide-1 mediates the therapeutic actions of DPP-IV inhibitors. *Diabetologia* 48:612–615. <https://doi.org/10.1007/s00125-005-1705-7>

59. Knop FK, Vilsbøll T, Højberg P V., et al (2007) Reduced incretin effect in type 2 diabetes: Cause or consequence of the diabetic state? *Diabetes* 56:1951–1959. <https://doi.org/10.2337/db07-0100>

60. Seino Y, Fukushima M, Yabe D (2010) GIP and GLP-1, the two incretin hormones: Similarities and differences. *J. Diabetes Investig.* 1:8–23. <https://doi.org/10.1111/j.2040-1124.2010.00022.x>

61. Fu Z, R. Gilbert E, Liu D (2012) Regulation of insulin synthesis and secretion and pancreatic beta-cell dysfunction in diabetes. *Curr Diabetes Rev* 9:25–53. <https://doi.org/10.2174/15733998130104>
62. Holz GG (2004) Epac: A new cAMP-binding protein in support of glucagon-like peptide-1 receptor-mediated signal transduction in the pancreatic  $\beta$ -cell. *Diabetes* 53:5–13. <https://doi.org/10.2337/diabetes.53.1.5>
63. Tengholm A (2012) Cyclic AMP dynamics in the pancreatic  $\beta$ -cell. *Ups. J. Med. Sci.* 117:355–369. <https://doi.org/10.3109/03009734.2012.724732>
64. Tomas A, Jones B, Leech C (2020) New insights into beta-cell GLP-1 receptor and cAMP signaling. *J. Mol. Biol.* 432:1347–1366. <https://doi.org/10.1016/j.jmb.2019.08.009>
65. Ackeifi C, Wang P, Karakose E, et al (2020) GLP-1 receptor agonists synergize with DYRK1A inhibitors to potentiate functional human  $\beta$  cell regeneration. *Sci Transl Med* 12:eaaw9996. <https://doi.org/10.1126/scitranslmed.aaw9996>
66. Nugent C, Prins JB, Whitehead JP, et al (2001) Potentiation of glucose uptake in 3T3-L1 adipocytes by PPAR $\gamma$  agonists is maintained in cells expressing a PPAR $\gamma$  dominant-negative mutant: Evidence for selectivity in the downstream responses to PPAR $\gamma$  activation. *Mol Endocrinol* 15:1729–1738. <https://doi.org/10.1210/mend.15.10.0715>
67. DeFronzo RA, Tripathy D (2009) Skeletal muscle insulin resistance is the primary defect in type 2 diabetes. *Diabetes Care* 32:S157-163. <https://doi.org/10.2337/dc09-S302>

68. Guilherme A, Virbasius J V., Puri V, Czech MP (2008) Adipocyte dysfunctions linking obesity to insulin resistance and type 2 diabetes. *Nat. Rev. Mol. Cell Biol.* 9:367–377. <https://doi.org/10.1038/nrm2391>
69. Basu A, Basu R, Shah P, et al (2001) Type 2 diabetes impairs splanchnic uptake of glucose but does not alter intestinal glucose absorption during enteral glucose feeding: Additional evidence for a defect in hepatic glucokinase activity. *Diabetes* 50:1351–1362. <https://doi.org/10.2337/diabetes.50.6.1351>
70. Dubreuil P, Fulcrand P, Rodriguez M, et al (1990) ACE-like hydrolysis of gastrin analogs and CCK-8 by fundic mucosal cells of different species with release of the amidated C-terminal dipeptide. *Biochim Biophys Acta Protein Struct Mol* 1039:171–176. [https://doi.org/10.1016/0167-4838\(90\)90182-F](https://doi.org/10.1016/0167-4838(90)90182-F)
71. Lafferty RA, Flatt PR, Irwin N (2018) C-terminal degradation of PYY peptides in plasma abolishes effects on satiety and beta-cell function. *Biochem Pharmacol* 158:95–102. <https://doi.org/10.1016/j.bcp.2018.10.004>
72. Fu Z, R. Gilbert E, Liu D (2012) Regulation of insulin synthesis and secretion and pancreatic beta-cell dysfunction in diabetes. *Curr Diabetes Rev* 9:25–53. <https://doi.org/10.2174/15733998130104>
73. Omar BA, Liehua L, Yamada Y, et al (2014) Dipeptidyl peptidase 4 (DPP-4) is expressed in mouse and human islets and its activity is decreased in human islets from individuals with type 2 diabetes. *Diabetologia* 57:1876–1883. <https://doi.org/10.1007/s00125-014-3299-4>
74. Bugliani M, Syed F, Paula FMM, et al (2018) DPP-4 is expressed in human pancreatic beta cells and its direct inhibition improves beta cell function and survival in type 2

1        diabetes.            Mol            Cell            Endocrinol            473:186–193.

2        <https://doi.org/10.1016/j.mce.2018.01.019>

3

4

## Figure Legends

**Figure 1.** (A-J) Effects of PPPH (0.0195 – 2.5 mg/ml) on insulin release from clonal pancreatic BRIN-BD11 beta-cells at basal, 5.6 mM, glucose concentration. (K) Additionally, effects of a fixed (2.5 mg/ml) concentration of various protein hydrolysate on cell viability were also investigated. Values are mean  $\pm$  SEM (n=8). \* $p$ <0.05, \*\* $p$ <0.01, \*\*\* $p$ <0.001 compared to control 5.6 mM glucose (A-K). Con: aqueous/alkaline control, AF: Alcalase/Flavourzyme, Alc: Alcalase, Brom: Bromelain, Prom: Promod.

**Figure 2.** (A-J) Effects of PPPH (0.0195–2.5 mg/mL) on insulin release from clonal pancreatic BRIN-BD11 beta-cells at elevated, 16.7 mM, glucose concentration. (K) Additionally, effects of a fixed (2.5 mg/ml) concentration of various protein hydrolysate on cell viability were also investigated (K). Values are mean  $\pm$  SEM (n=8). \*\* $p$ <0.01, \*\*\* $p$ <0.001 compared to control 16.7 mM glucose (A-K). Con: aqueous/alkaline control, AF: Alcalase/Flavourzyme, Alc: Alcalase, Brom: Bromelain, Prom: Promod.

**Figure 3.** Effects of fixed concentration (2.5 mg/mL) of PPPH on the incretin effect through (A) GLP-1 release from GLUTag cells and (B) GIP release from STC-1  $\alpha$ -cells. Values are mean  $\pm$  SEM (n=4).  $p$ <0.05, \*\* $p$ <0.01, \*\*\* $p$ <0.001 compared to 2 mM glucose controls (A,B).  $^{\Delta}p$ <0.05,  $^{\Delta\Delta}p$ <0.01,  $^{\Delta\Delta\Delta}p$ <0.001 compared to the appropriate, non-SGID hydrolysate (A,B). Con: aqueous/alkaline control, AF: Alcalase/Flavourzyme, Alc: Alcalase, Brom: Bromelain, Prom: Promod.

**Figure 4.** Mechanistic effects of coincubation with fixed concentration (2.5 mg/ml) of PPPH following 20 min incubation with BRIN-BD11 cells. (A) PPPH's and SGID equivalents, were tested for influence on cAMP concentration, (B) while only the Alcalase/Flavourzyme PPPH (AF) was employed for the study of intracellular calcium mobilisation and (D) membrane potential. Respective AUC values are also provided (C,E). Values are mean  $\pm$  SEM (n=3).

\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  compared to 5.6 mM glucose control (A,C,E).  $\Delta\Delta p < 0.001$  compared to the 10 mM alanine positive control (C,E). Con: aqueous/alkaline control, AF: Alcalase/Flavourzyme, Alc: Alcalase, Brom: Bromelain, Prom: Promod.

**Figure 5.** Effects of PPPH (2.5 mg/ml) on glucose uptake individually, or in combination with insulin (1 nM), in trans-differentiated 3T3-L1 adipocyte cells. Responsiveness of adipocyte cells was assessed via incubation with either apigenin (50  $\mu$ M), insulin (1 nM or 100 nM) (A). Adipocyte cells were further incubated with either Con/Con-SGID (B), Alc/Alc-SGID (C), AF/AF-SGID (D), Brom/Brom-SGID (E) or Prom/Prom-SGID (F) using fixed concentration of hydrolysate (2.5 mg/ml) for 1 h with 3 mM fluorescent glucose (2-NBDG). Values are mean  $\pm$  SEM (n=3). \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  compared to respective glucose control. Con: aqueous/alkaline control, AF: Alcalase/Flavourzyme, Alc: Alcalase, Brom: bromelain, Prom: Promod.

**Figure 6.** (A,B) Effects of a fixed dose (100 mg/kg/bw) of Alcalase/Flavourzyme PPPH (AF) on glucose tolerance in lean, overnight-fasted (16 h) NIH Swiss mice and (C) acute food intake in lean, diet-restricted HsD:Ola T0 mice. Animals (n=8) received glucose (18.8 mmol/kg/bw) alone or in combination with hydrolysate (A) or saline (0.9% NaCl) alone or in combination with hydrolysate via oral gavage (C). Measurements were taken at regular intervals as indicated



- 1 (A,C). Values are mean  $\pm$  SEM (n=8). \* $p$ <0.05, \*\* $p$ <0.01, \*\*\* $p$ <0.001 compared to glucose
- 2 only (A,B) or saline only (C) controls.

Figure 1

Figure 1




Figure 2

Figure 2





Figure 3

Figure 3

A



B




Figure 4

A






Figure 4


B



C



D



E




Figure 5









Figure 6






Figure 6



**Table 1.** DPP-4 inhibitory activity of PPPH's and their SGID equivalents

Proteolytic activity	IC <sub>50</sub> value (mg/ml)	
	PPPH	PPPH-SGID
Control (none)	1.91 ± 0.10 <sup>d</sup>	1.09 ± 0.06 <sup>bc</sup> *
Alcalase + Flavourzyme	0.70 ± 0.02 <sup>a</sup>	1.00 ± 0.03 <sup>bc</sup> *
Alcalase	0.94 ± 0.10 <sup>b</sup>	1.14 ± 0.05 <sup>c</sup> *
Bromelain	1.34 ± 0.05 <sup>c</sup>	0.95 ± 0.08 <sup>b</sup> *
Promod	1.23 ± 0.05 <sup>c</sup>	0.78 ± 0.02 <sup>a</sup> *

Mean ± SD (n=3), IC<sub>50</sub>: inhibitory concentration that inhibits enzyme activity by 50%. \* indicates a significant difference (*p*< 0.05) in IC<sub>50</sub> values following SGID.





[Click here to access/download](#)

**Electronic Supplementary Material**  
Supplementary Figures Eur J Nut 01-02-2021.pptx

